A CERTAIN SUBCLASS OF MEROMORPHICALLY STARLIKE FUNCTIONS WITH ALTERNATING COEFFICIENTS

M.K. Aouf, H.M. Hossen and A.Y. Lashin

Abstract. Coefficient inequalities and distortion theorems are obtained for certain starlike functions with alternating coefficients. Further class preserving integral operators are obtained.

MSC 2000. 30C45.

Key words. Analytic alternating coefficients, starlike functions.

REFERENCES

- [1] AOUF, M.K., A certain subclass of meromorphic starlike functions with positive coefficients, Rend. Mat. Roma (7) 9, (1989), 225-235.
- [2] AOUF. M.K. AND DARWISH, H.E., Meromorphic starlike univalent functions with alternating coefficients, Utilitas Math. 47 (1995), 137-144.
- [3] Cluine, J., On meromorphic schlicht functions, J. London Math. Soc. 34 (1959), 215-216.
- [4] MILLER, J.E., Convex meromorphic mapping and related functions, Proc. Amer. Math. Soc. 25 (1970), 220-228.
- [5] MOGRA, M.L., REDDY, T.R. AND JUNEJA, O.P., Meromorphic univalent functions with positive coefficients, Bull. Austral. Math. Soc. 32 (1985), 161-176.
- [6] Pommerenke, C., On meromorphic starlike functions with positive coefficients, Pacific J. Math. 13 (1963), 221-235.
- [7] URALEGADDI, B.A. AND GANIGI, M.D., Meromorphic starlike functions with alternating coefficients, Rend. Mat. Roma (7) 11, (1991),441-446.
- [8] URALEGADDI, B.A. AND SOMANTHA, C., New criteria for meromorphic starlike univalent functions with positive coefficients, Bull. Astral. Math. Soc. 43 (1991),137-140.

Received: May 21, 1999

Department of Mathematics

Faculty of Science

Mansoura University

Mansoura, Egypt

e-mail: sinfac@mum.muns.eun.eg