## SUFFICIENT CONDITIONS FOR QUASICONFORMAL EXTENSIONS IN SEVERAL COMPLEX VARIABLES

## PAULA CURT

**Abstract.** Let B be the unit ball in  $\mathbb{C}^n$  with respect to the euclidian norm. In this paper we will give a sufficient condition such that a holomorphic mapping defined on B can be extended to a quasiconformal homeomorphism of  $\mathbb{R}^{2n}$  onto itself.

MSC 2000. 30C65.

**Key words.** Holomorphic maps, subordination.

## REFERENCES

- [1] BECKER, J., Löewnersche Differentialgleichung und quasikonform fortsetzbare schlichte Funktionen, J. Reine Angew. Math., 255 (1972), 23–43.
- [2] BOJARSKI, B. and IWANIEC, T., Analytical foundations of the theory of quasiconformal mappings in  $\mathbb{R}^n$ , Ann. Acad. Sci. Fenn. Ser A Math., 8 (1983), 257–324.
- [3] Chuaqui, M., Applications of subordination chains to starlike mappings in C<sup>n</sup>, Pacif. J. Math., 168 (1995), 33–48.
- [4] Curt, P., A generalization in n-dimensional complex space of Ahlfors' and Becker's criterion for univalence, Studia Univ. Babeş-Bolyai, Math., 39 (1994), 31–38.
- [5] Curt, P., Subordination chains and quasiconformal extensions in several variables, to appear.
- [6] Curt, P. and Kohr, G., Subordination chains and Loewner differential equation in several complex variables, to appear.
- [7] HAMADA, H. and KOHR, G., The growth theorem and quasiconformal extension of strongly spirallike mappings of type α, Complex variables, 44 (2001), 281–297.
- [8] HAMADA, H. and KOHR, G., Quasiconformal extension of strongly starlike mappings on the unit ball in  $\mathbb{C}^n$ , Mathematica (Cluj), to appear.
- [9] PFALTZGRAFF, J.A., Subordination chains and quasiconformal extension of holomorphic maps in  $\mathbb{C}^n$ , Ann. Acad. Sci. Fennicae, Ser. A, Mathematica, 1 (1975), 13–25.

Received January 30, 2002

Faculty of Mathematics and Computer Science
"Babeş-Bolyai" University
Str. M. Kogălniceanu 1
RO-400084 Cluj-Napoca, România
E-mail: paula@math.ubbcluj.ro