

BOUNDS ON THE COEFFICIENTS OF CERTAIN ANALYTIC
AND UNIVALENT FUNCTIONS

K. O. BABALOLA

Abstract. For the real number $\alpha > 1$, we use a technique due to Nehari and Netanyahu and an application of certain integral iteration of Caratheodory functions to find the best-possible upper bounds on the coefficients of functions of the class $T_n^\alpha(\beta)$ introduced in [4] by Opoola.

MSC 2000. 30C45.

Key words. Coefficient bounds, analytic and univalent functions.

REFERENCES

- [1] BABALOLA, K. O. and OPOOLA, T. O., *Iterated integral transforms of Caratheodory functions and their applications to analytic and univalent functions*, Tamkang J. Math., **37** (4) (2006), 355–366.
- [2] BABALOLA, K. O. and OPOOLA, T. O., *On the coefficients of certain analytic and univalent functions*, Advances in Inequalities for Series, Nova Science Publishers (<http://www.novapublishers.com>) (2006), 5–17. (Edited by S. S. Dragomir and A. Sofo).
- [3] NEHARI, Z. and NETANYAHU, E., *On the coefficients of meromorphic schlicht functions*, Proc. Amer. Math. Soc. **8** (1) (1957), 15–23.
- [4] OPOOLA, T. O., *On a new subclass of univalent functions*, Matematica (Cluj) **36**, 59 (2)(1994), 195–200.
- [5] OPOOLA, T. O., BABALOLA, K. O., FADIPE-JOSEPH O. A. and RAUF, K., *On the coefficient bounds of a subclass of univalent functions*, J. Nig. Math. Soc. **24** (2004), 87–92.
- [6] SINGH, R., *On Bazilevic functions*, Proc. Amer. Math. Soc. **38** (1973), 261–271.

Received March 28, 2007

Department of Mathematics

University of Ilorin

Ilorin, Nigeria

E-mail: abuuabdilqayyuum@gmail.com