

KRASNOSELSKII-TYPE THEOREMS FOR MULTIVALUED OPERATORS

MONICA BORICEANU

Department of Applied Mathematics, Babeş-Bolyai University
1, Kogălniceanu Str., 400084, Cluj-Napoca, Romania
E-mail: bmonica@math.ubbcluj.ro

Abstract. The aim of this paper is to present some fixed point theorems of Krasnoselskii-type for the sum of two multivalued operators.

Key Words and Phrases: Fixed point, multivalued operator, densifying operator, contraction.

2000 Mathematics Subject Classification: 47H10, 54H25.

REFERENCES

- [1] Teck-Cheong Lim, *On characterizations of Meir-Keeler contractive maps*, *Nonlinear Analysis* 46(2001), 113-120.
- [2] M. Furi, A. Vignoli, *Fixed points for densifying mappings*, *Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur.*, (8) **47**(1970), 465-467.
- [3] M.Z. Nashed, J.S. Wong, *Some variants of a fixed point theorem of Krasnoselskii and applications to nonlinear integral equations*, *J. Math. Mech.*, **18**(1969), 767-777.
- [4] A. Petruşel, *Operatorial Inclusions*, House of the Book of Science, 2002.
- [5] S. Reich, *Fixed point of contractive functions*, *Boll. U.M.I.*, **5**(1972), 26-42.
- [6] L. Rybinski, *Multivalued contraction with parameter*, *Ann. Polon. Math.*, **45**(1985), 275-282.
- [7] R. Wegrzyk, *Fixed point theorems for multifunctions and their applications to functional equations*, *Diss. Math.*, **201**(1982).

Received: November 21, 2007; Accepted: February 6, 2008.

This paper was presented at the International Conference on Nonlinear Operators, Differential Equations and Applications held in Cluj-Napoca (Romania) from July 4 to July 8, 2007.