

TREND CONSTANTS FOR LIPSCHITZ MAPPINGS

KRZYSZTOF BOLIBOK AND KAZIMIERZ GOEBEL

Institute of Mathematics, Maria Curie-Skłodowska University
20-031 Lublin, Poland

E-mails: bolibok@hektor.umcs.lublin.pl, goebel@hektor.umcs.lublin.pl

Abstract. Lipschitz mappings are naturally classified by the value of their Lipschitz constant. Here we propose to use two other *trend constants* which allow a more subtle classification of such mappings. Basic facts, examples and applications to metric fixed point theory are shown.

Key Words and Phrases: Lipschitz mappings, nonexpansive mappings, fixed points.

2010 Mathematics Subject Classification: 47H10, 54H25.

REFERENCES

- [1] F.E. Browder, *Nonexpansive nonlinear operators in a Banach space*, Proc. Nat. Acad. Sci. USA, **56**(1965), 1041-1044.
- [2] R.E. Bruck, *Properties of fixed point sets of nonexpansive mappings*, Trans. Amer. Math. Soc., **179**(1973), 251-262.
- [3] K. Goebel, *Concise Course on Fixed Point Theorems*, Yokohama Publishers, 2002.
- [4] K. Goebel, S. Prus, *Shapes and sizes of fixed point sets of nonexpansive mappings*, Nonlinear Analysis and Convex Analysis I, Yokohama Publishers (2013) 75-98.
- [5] K. Goebel, W.A. Kirk, *Topics in Metric Fixed Point Theory*, Cambridge University Press, 1990.
- [6] D. Göhde, *Zum princip der kontraktiven Abbildung*, Math. Nachr., **30**(1965), 251-258.
- [7] W.A. Kirk, *A fixed point theorem for mappings which do not increase distances*, Amer. Math. Monthly, **72**(1965), 1004-1006.
- [8] W.A. Kirk, B. Sims, *Handbook on Metric Fixed Point Theory*, Kluwer Academic Publ., 2001.

Received: January 30, 2013; Accepted: March 22, 2014.